

John A. Selling, MD
Mary F. Chan, MD
Waverley Surgery Center
400 Forest Avenue
Palo Alto, CA 94301

Doctor's phone number: 650-498-6690

Waverley Phone: 650-324-0600

Waverley Fax: 650-289-1620

PROCEDURE: Colonoscopy

FACILITY: Waverley Surgery Center

PATIENT: _____

DATE OF EXAM: _____

ARRIVAL TIME: _____

PLEASE READ THESE INSTRUCTIONS AT LEAST ONE WEEK PRIOR TO YOUR COLONOSCOPY. Following this schedule will prepare you for a safe and convenient experience. The cleaner your colon is, the better the quality of the exam.

Five days before colonoscopy:

- Please stop taking any form of aspirin and Plavix. Tylenol (acetaminophen) may be taken for pain or headache.
- If you have kidney disease, please notify Dr. Selling/Chan's office as we will give you a different prep.
- If you have diabetes, please check with your primary care doctor for instructions on how to manage your diabetes medicines on the day before and the day of your colonoscopy.
- If you have an internal defibrillator, we need to do your exam at Stanford Medical Center. Call Dr. Selling/Chan's office to reschedule. Pacemakers are fine.
- Take the prescription for Osmoprep to your pharmacy and purchase it. You may also buy Vitamin A and D ointment or petroleum jelly to apply to your perianal area during the prep if desired.
- Arrange for a ride home for the day of your exam. This person should be known to you and by law, has to be at least 18 years of age. If you are disabled or anticipate difficulty with mobility in a semi-sedated state, it would be advantageous to have this person stay with you for a few hours after arriving home. Taxis and trains are acceptable only if you are accompanied by an adult companion. If needed, an acceptable transportation service is Older Adults Care Management (650)329-1411

Three days before colonoscopy:

- Stop taking Coumadin (warfarin). If you have an artificial heart valve, you should discuss an alternative anticoagulant with your cardiologist or primary care doctor. Cutting your usual dose of Coumadin in half for 3 days is an alternative in this case if acceptable to your cardiologist.
- Avoid eating nuts, seeds, popcorn, grapes, peas, onions, beans, and tomatoes.

Two days before colonoscopy:

- Please stop all anti-inflammatory medications (Ibuprophen, Advil, Motrin, Mobic, Indomethacin, Aleve, Datriil, Naproxen, Naprosyn, Nuprin).

One day before colonoscopy:

- THESE INSTRUCTIONS ARE IMPORTANT. YOUR COLON MUST BE CLEAN.
- Start drinking clear liquids in the morning and continue throughout the day. Do **not** ingest solid foods, milk or milk products. Clear liquids include all of the following that are not red or purple; fruit juices (apple, white grape, lemonade), water, clear broth or boullion, black coffee or tea, Gatorade, carbonated and non-carbonated soft drinks, Jello and popsicles. Other than as mentioned above, you may take your usual medications today.
- Start the first regimen of Osmoprep at 5pm the afternoon prior to your exam. (You may start earlier or later than this, but it is important that there are at least 6 hours between the starting times of the 2 regimens of Osmoprep.) The first regimen consists of one dose (4 tablets with 8 oz. of any clear liquid) every 15 minutes for a total of 5 doses (20 tablets and 40 oz. of clear liquid). This prep may work within 30 minutes, but may take as long as 4 hours. Remain close to the bathroom, as multiple bowel movements will occur. Bloating, nausea and/or chilling may occur, but will improve when bowel movements begin. If nausea is significant, stop for 30 minutes, then resume drinking. Consider using petroleum jelly around the anus before starting and during the prep. Make a conscious effort to drink as much as you can before, during and after the prep to prevent dehydration.
- Repeat the process with the second regimen of Osmoprep at 11pm (or at least 6 hours after starting the first regimen). The second regimen consists of one dose (4 tablets with 8 oz. of clear liquid) every 15 minutes for a total of 3 doses (12 tablets and 24 oz. of clear liquid). It is important that you drink at least the minimum of required liquid after each dose because the prep works by washing your colon clean. (The second regimen is required even if your bowel movements are liquid after the first regimen and even if you are small in weight or stature.)
- If your appointment arrival time is after 10:00am, you have the option of taking the 2nd regimen of Osmoprep at 6:00am on the same day as your exam. Administer the regimen as directed above. **All liquids must be finished 2 hours before arrival.**

Day of colonoscopy:

- Take your usual medications (not mentioned above) at least 2 hours before arrival. Do not take morning blood pressure medications unless your arrival time is after noon.
- **You may have only clear liquids until two hours before your arrival time at the surgical center, nothing thereafter.**
- Please arrive on time. Parking is underneath the building. Take the elevator to the first floor and go to the Reception desk. You will be at the surgical center for 2 hours.
- Wear loose, comfortable clothing.
- Bring your photo ID and your insurance card.
- A nurse will check you in and insert an IV catheter into a vein.

- During the exam your cardiovascular status will be monitored. Supplemental oxygen will be given via a nasal cannula. An IV medication will be given which will make you sleepy and relaxed. You will be comfortable lying on your left side during the exam. The colonoscope will be passed through your entire large intestine. If necessary, small biopsies may be taken and, if present, polyps (small growths) may be removed. You will feel no discomfort if this is done.

After the colonoscopy:

- You will be observed until you adequately recover from the effects of the medication. If you desire, the doctor may discuss your findings with your ride if he/she is available. Discharge instructions will be given to you and your ride in the recovery area. You cannot drive until the following morning. You may eat a regular diet and take your usual medications when you arrive home unless otherwise directed at discharge.

Are there any complications from colonoscopy and polypectomy?

- Colonoscopy and polypectomy are associated with very low risk when performed by doctors with special training and experience in these endoscopic procedures. One possible complication is perforation, a tear through the wall of the bowel. Another complication is bleeding which may occur from the site of biopsy or polyp removal. It is usually minor and stops on its own or by cauterization (application of electrical current) during the exam. Rarely, surgery is necessary. A third complication is a localized irritation of the vein in your arm that may occur at the site of medication injection. A tender lump could develop and may remain for several weeks to months. Elevation of the arm and local application of heat speeds recovery. A fourth possible complication may occur if the doctor is unable to visualize an abnormality due to poor colon preparation. It is important that the colon be clean to get the best visualization.

Causes for concern warranting a call to your doctor:

- Rectal bleeding greater than one quarter cupful, or a lesser amount lasting more than one to two days.
- Increasing abdominal pain.
- Nausea or vomiting.
- Fever or chills.

Please note:

- If you have any questions about your exam, please feel free to discuss them with Dr. Selling/Chan or the nurse at 650-498-6690.
- You will be responsible for both a facility fee from Waverley Surgery Center and the physician's professional fee. If a biopsy is done, there will be a separate charge from Stanford Pathology. If you need to cancel your appointment for any reason, please call the nurse at least 48 hours in advance. There is a \$50.00 no-show charge per Menlo Medical Clinic management. Thank you for trusting us with your care.